

# Tafluprost INN 0.0015% Sterile Ophthalmic Solution

#### Composition:

Taflan Sterile Ophthalmic Solution: Each ml contains Tafluprost INN 0.015 mg.

#### Preservative

Sodium Perborate BP 0.01%

### Pharmacology:

Tafluprost, a prostaglandin analog is a selective FP prostanoid receptor agonist which is believed to reduce intraocular pressure by increasing uveoscleral outflow of aqueous humor.

#### Indications:

**Taflan** is indicated for reducing elevated intraocular pressure in patients with open-angle glaucoma or ocular hypertension.

# Dosage & administration:

Instill one drop in the affected eye(s) once daily in the evening.

### Contraindications:

Tafluprost ophthalmic solution is contraindicated in patients who are hypersensitive to any component of this product.

#### Side effects

The most frequently reported treatment-related side effect is ocular hyperemia. Rare side effects observed include eye irritation and blurred vision.

# Precautions:

- For ophthalmic use only.
- To avoid possible contamination of the drops, do not touch the dropper tip or to any surface.
- Should be used with caution in patients with active intraocular inflammation (e.g., iritis/uveitis).
- If more than one topical ophthalmic drug is being used, the drugs should be administered at least five (5) minutes between applications.

# Use in special group:

**Use in pregnancy:** Pregnancy category C. There are no adequate and well-controlled studies in pregnant women. Tafluprost ophthalmic solution should be used during pregnancy unless the potential benefit justifies the potential risk to the fetus.

**Use in lactation:** It is not known whether Tafluprost is excreted in human milk. Caution should be exercised while giving this ophthalmic solution to a nursing mother.

Use in children: Use in pediatric patients is not recommended.

Use in elderly patients: No overall differences in safety and effectiveness have been observed between elderly and other adult patients.

# Storage

- Keep unopened container at 2 8°C. After opening keep the container in a cool & dry place below 25°C.
- Do not use longer than 30 days after first opening.
- Keep out of the reach of children.

# Packing

Taflan Sterile Ophthalmic Solution: Each plastic dropper bottle contains 3 ml sterile ophthalmic solution.



