

# ***Nevan & Nevan TS***

Nepafenac INN  
Sterile Ophthalmic Suspension

**Composition:**

**Nevan Sterile Ophthalmic Suspension:** Each ml contains Nepafenac INN 1 mg.  
**Nevan TS Sterile Ophthalmic Suspension:** Each ml contains Nepafenac INN 3 mg.

**Preservative:**

Sodium Perborate 0.01%.

**Vehicle:**

**Nevan Sterile Ophthalmic Suspension:** Carbomer 980 USP 0.4%.  
**Nevan TS Sterile Ophthalmic Suspension:** Carbomer 980.

**Pharmacology:**

Nepafenac is a nonsteroidal anti-inflammatory prodrug (NSAID). After instillation in the eye, it penetrates the cornea and is converted by ocular tissue hydrolase to Amfenac, a potent nonsteroidal anti-inflammatory drug. Amfenac is thought to inhibit the action of cyclooxygenase enzyme. This enzyme is required for prostaglandin synthesis.

**Indications:**

**Nevan Sterile Ophthalmic Suspension-**

- The treatment of post-operative ocular pain and inflammation including cataract surgery
- Inhibition of surgery induced miosis and
- Prevention of post-operative cystoid macular edema (CME).

**Nevan TS Sterile Ophthalmic Suspension-**

- Post-operative pain and inflammation associated with cataract surgery.

**Dosage & administration:**

**Nevan Sterile Ophthalmic Suspension:** For post-operative pain & inflammation: Instill 1 drop 3 times daily 1 day prior to cataract surgery and continued on the day of surgery and through the first 2 weeks of the post-operative period.

For surgery induced miosis: Instill 1 drop 3 times daily 1 day before surgery & on the day of surgery.

For prevention of post-operative CME: Instill 1 drop 3 times daily 1 day before surgery and continued on the day of surgery and through the first 6 weeks of the post-operative period.

**Nevan TS Sterile Ophthalmic Suspension:** For post-operative pain & inflammation: Instill 1 drop once daily 1 day prior to cataract surgery and continued on the day of surgery and through the first 2 weeks of the post-operative period. An additional drop should be administered 30 to 120 minutes prior to surgery.

**Contraindications:**

Contraindicated in patients with previously demonstrated hypersensitivity to any of the ingredients in the formulation or to other NSAIDs.

**Side effects:**

Reported side effects are foreign body sensation, lid margin crusting, ocular discomfort, ocular hyperemia etc.

**Precautions:**

Nepafenac should be used with caution in patients with known bleeding tendencies or who are receiving medications which may prolong bleeding time.

**Use in special group:**

**Use in pregnancy:** There are no adequate and well-controlled studies in pregnant women. Nepafenac should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

**Use in lactation:**

It is not known whether this drug is excreted in human milk. So, caution should be exercised when Nepafenac ophthalmic suspension is administered to a nursing mother.

**Use in children:**

Safety and effectiveness in pediatric patients below 18 years of age have not been established.

**Use in elderly patients:** No overall differences in safety or effectiveness have been observed between elderly and younger patients.

**Storage:**

- Store below 30°C, protect from light.
- Do not use longer than 30 days after first opening.
- Keep out of the reach of children.

**Packing:**

**Nevan Sterile Ophthalmic Suspension:** Each plastic dropper bottle contains 5 ml sterile ophthalmic suspension.

**Nevan TS Sterile Ophthalmic Suspension:** Each plastic dropper bottle contains 5 ml sterile ophthalmic suspension.



Manufactured by:  
**ARISTOPHARMA LTD.**  
SHAMPUR-DHAKA-BANGLADESH

20002372/05