

Matropin

Homatropine Hydrobromide USP 2%
Eye Drops

Composition:

Matropin Eye Drops: Each ml contains Homatropine Hydrobromide USP 20 mg.

Preservative:

Sodium Perborate BP 0.01%.

Pharmacology:

Matropin contains Homatropine which blocks the action of certain parasympathetic nerves and cholinergic drugs. The effect is exerted in 15 to 30 minutes and passes off in 12 to 24 hours.

Indications:

Matropin Eye Drop is indicated for use as a mydriatic and cycloplegic.

Dosage & administration:

For refraction: Instill two drops in the eye(s) and repeat 2-3 times at 5-10 minute intervals.

For therapy: Instill 1-2 drops in the eye(s) every 3-4 hours.

Contraindications:

Contraindicated in patients with closed-angle glaucoma or with a narrow angle between the iris and the cornea.

Side effects:

Hypersensitivity may occur as conjunctivitis.

Precautions:

Should be used with caution in patients with prostatic enlargement and in patients suffering from paralytic ileus or pyloric stenosis.

Use in special group:

Use in pregnancy: There are no adequate and well-controlled studies in pregnant women. Matropin should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Use in lactation: It is not known whether Homatropine is excreted in human milk. Caution should be exercised when Matropin is administered to a nursing mother.

Use in children: Should not be used in children below the age of 3 months.

Storage:

- Store in a cool & dry place, protect from light.
- Do not use longer than 30 days after first opening.
- Keep out of the reach of children.

Packing:

Matropin Eye Drops: Each LDPE dropper bottle contains 10 ml sterile & pyrogen free eye drops.

20002306/02



Manufactured by:
ARISTOPHARMA LTD.
SHAMPUR-DHAKA-BANGLADESH