

# NCL

Sodium Chloride 5%  
(Hypertonic Solution)

**Composition:**

**NCL Eye Drops:** Each ml solution contains Sodium Chloride BP 50 mg.

**Preservative :** Sodium perborate.

**Indications:**

NCL Eye Drops is indicated for the temporary relief of corneal edema. It may also be used during eye exams.

**Dosage & administration:**

Instill 1 drop in the affected eye (s) every 3 or 4 hours or as directed by a physician.

**Contraindications:**

Contraindicated in patients who develop allergic reactions to this medicine.

**Side effects:**

Vision may be temporarily blurred for a period after applying the drops. Burning or irritation of the eye may occur temporarily when first applied. Other side effects are redness, pain, swelling in or around the eyes, sensitivity to light and headache.

**Precautions:**

Precaution should be taken if patients have history of eye problems or allergies.

**Use in special group:**

**Use in pregnancy & lactation:** As with any medication, this should be used cautiously during pregnancy or while breast-feeding.

**Use in children:** Although there is no specific information comparing use of hypertonic sodium chloride in children with use in other age groups, this medicine is not expected to cause different side effects or problems in children than it does in adults.

**Use in elderly patients:** There is no specific information comparing use of hypertonic sodium chloride in the elderly with use in other age groups.

**Storage:**

- Store below 30° C temperature in a dry place, protect from light.
- Keep out of the reach of children.
- Discard this if the solution turns color, cloudy or contains particles.

**Packing:**

**NCL Eye Drops:** Each LDPE dropper bottle contains 5 ml Sterile Eye Drops.

20005744/00



Manufactured by:  
**ARISTOPHARMA LTD.**  
GAZIPUR, BANGLADESH