

Olpadin

Olopatadine Hydrochloride USP
Eye Drops

Composition:

Olpadin Eye Drops: Each ml contains Olopatadine Hydrochloride USP equivalent to Olopatadine 1 mg.

Olpadin DS Eye Drops: Each ml contains Olopatadine Hydrochloride USP equivalent to Olopatadine 2 mg.

Olpadin 7 Eye Drops: Each ml contains Olopatadine Hydrochloride USP equivalent to Olopatadine 7 mg.

Preservative: Sodium Perborate BP 0.01%.

Vehicle:

Olpadin and Olpadin DS Eye Drops: Polyvinyl Alcohol USP 1.4%

Olpadin 7 Eye Drops: Povidone K-17

Pharmacology:

Olopatadine selectively blocks histamine H₁ receptor and stabilizes mast cells. Thus, it prevents histamine binding to its receptors and inhibits mast cell degranulation & histamine release. It also decreases chemotaxis and inhibits the activation of eosinophil.

Indications:

Olpadin, Olpadin DS & Olpadin 7 are indicated for the treatment of ocular itching associated with allergic conjunctivitis.

Dosage and administration:

Olpadin: Instill 1 drop in the affected eye(s) twice daily.

Olpadin DS: Instill 1 drop in the affected eye(s) once daily.

Olpadin 7: Instill 1 drop in the affected eye(s) once daily.

Contraindications:

Contraindicated in patients who are hypersensitive to Olopatadine Hydrochloride or any other component of the preparation.

Side effects:

Blurred vision, temporary burning & stinging, dry eye, foreign body sensation, hyperemia, dysgeusia.

Precautions:

- For ophthalmic use only
- To avoid possible contamination of the drops, do not touch the dropper tip or to any surface
- Do not wear a contact lens during the treatment
- If more than one topical ophthalmic drug is being used, the drugs should be administered at least (5) minutes interval between applications

Drug Interactions:

Olopatadine has low drug interaction potential as systemic levels of Olopatadine achieved after ocular dosing are negligible.

Use in special group:

Use in Pregnancy: There are no adequate and well-controlled studies in pregnant women. Olopatadine should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Use in Lactation: It is not known whether Olopatadine passes into breast milk. However, caution should be exercised when Olopatadine is administered to a nursing mother.

Use in Pediatric:

Safety and effectiveness in children for Olopatadine 0.1% is below the age of 3 years and Olopatadine 0.2% & 0.7% is below 2 years have not been established.

Use in Geriatric: No overall differences in safety and effectiveness have been observed between elderly and younger patients.

Storage:

- **Olpadin & Olpadin DS Eye Drops:** Store at 4°-25°C.
- **Olpadin 7 Eye Drops:** Store below 25°C.
- Do not use longer than 30 days after first opening.
- Keep out of the reach of the children.

Packing:

Olpadin Eye Drops: Each LDPE dropper bottle contains 5 ml Sterile & Pyrogen free eye drops.

Olpadin DS Eye Drops: Each LDPE dropper bottle contains 5 ml Sterile & Pyrogen free eye drops.

Olpadin 7 Eye Drops: Each LDPE dropper bottle contains 5 ml Sterile & Pyrogen free eye drops.



Manufactured by:
ARISTOPHARMA LTD.
Shampur-Kadamtai I/A, Dhaka-Bangladesh

20002402/08