

A-MYCIN[®] Lotion

Erythromycin 3% w/v

Composition :

Each ml Lotion contains Erythromycin Ethylsuccinate USP equivalent to Erythromycin 30 mg.

Pharmacology :

A-MYCIN[®] Lotion (Erythromycin) is a bacteriostatic macrolide antibiotic, but may be bactericidal in high concentrations. Although the mechanism by which topical Erythromycin acts in reducing inflammatory lesions of acne vulgaris is unknown, it is presumably due to its antibiotic action.

Indications :

For topical treatment of acne, pimples & bacterial skin infections susceptible to Erythromycin.

Dosage & Administration :

- Apply to the affected areas in the morning and evening.
- Before applying thoroughly wash with warm water and soap, rinse and pat dry all areas to be treated.
- Apply with applicator.
- Wash hands after use.

Contraindications :

Hypersensitivity to Erythromycin or to any component of the product.

Precautions :

- For external use only.
- Keep away from eyes, nose, mouth and other mucous membrane.

Side Effects :

Erythema, desquamation, burning sensation, eye irritation, tenderness, dryness, oily skin, etc.

Use in Pregnancy :

Safety for use during pregnancy has not been established. Use only when the potential benefits outweigh potential hazards to the foetus.

Use in Lactation :

Erythromycin is excreted in breast milk. Exercise caution when administering to a nursing mother.

Paediatric Use :

Safety and efficacy in children < 12 years of age have not been established.

Superinfections :

Use of antibiotics (especially prolonged or repeated therapy) may result in bacterial or fungal overgrowth of non-susceptible organisms. Such overgrowth may lead to a secondary infection. Take appropriate measures if superinfection occurs.

Drug Interaction :

Clindamycin interacts with Erythromycin.

Storage :

- Store below 30°C, keep in dry place & protect from light.
- Keep out of the reach of children.

Packing :

Bottle containing 25 ml lotion.



Manufactured by:

ARISTOPHARMA LTD.

Shampur-Kadamtali I/A, Dhaka-Bangladesh


How to use A-MYCIN[®] Lotion



Shake the bottle before use.




Open the bottle & discard the cap.



Assemble the reticulated cork on its mouth.



Wash, rinse & dry the affected areas, and apply the lotion smoothly.



After using close the bottle well with the protective cap.