

Clobesol SA

Clobetasol Propionate USP 0.05% & Salicylic Acid BP 3%

Ointment

Composition:

Clobesol SA Ointment: Each gram ointment contains Clobetasol Propionate USP 0.5 mg and Salicylic Acid BP 30 mg.

Pharmacology:

Clobetasol Propionate has both local anti-inflammatory and immunosuppressive activity. The drug blocks the effect of macrophage migration inhibitory factor and decreases the activation of plasminogen to plasmin. It inhibits phospholipase A₂ activity via formation of lipocortin which reduces the formation of prostaglandins and leukotrienes in the local tissues. Salicylic acid is a keratolytic agent. It encourages exfoliation of the skin and opening up of the plugged follicles, which helps re-establish the normal skin-cell replacement cycle.

Indications:

- Initial control of all forms of hyper acute eczema
- Chronic hyperkeratotic eczema of the hands and feet and patches of chronic lichen-simplex
- Chronic hyperkeratotic psoriasis
- Severe acute photosensitivity
- Acute contact dermatitis
- Hypertrophic lichen planus & lichen sclerosis
- Localized bullous disorders
- Keloid scarring
- Pretibial myxoedema, vitiligo, suppression of reactions after cryotherapy
- Seborrheic dermatitis of the scalp

Dosage & Administration:

Apply the ointment as thin layer to the affected area twice daily. **Clobesol SA** treatment should not be continued for more than 14 days.

Side Effects:

Clobetasol may cause thinning of epidermis, burning, itching, irritation etc. Salicylic acid may cause allergic reaction, hives etc.

Contraindications:

Hypersensitivity to clobetasol propionate and salicylic acid.

Precautions:

- Avoid contact with eyes, mouth, nose or mucus membrane.
- Avoid occlusive dressing and application on the large surface area.

Use in Special Group:

Use in pregnancy: This combination should be avoided in pregnant women.

Use in lactation: Mothers using large amounts of the drug should be aware of potential excretion in milk.

Paediatrics Use: This combination is contraindicated in patients below 12 years.

Drug Interactions:

No hazardous interactions have been found.

Storage:

- Store at 15°C-30°C temperature and dry place, protect from light.
- Keep out of the reach of children.
- Close the mouth of tube well after each application.
- Do not freeze.

Packing:

Clobesol SA Ointment: Each tube contains 10 g/30 g ointment.

20002105/02

Manufactured by:

 **ARISTOPHARMA LTD.**
SHAMPUR-DHAKA-BANGLADESH